

Correction

Correction: Prediction of Muscle Activities from Electrocorticograms in Primary Motor Cortex of Primates

The PLOS ONE Staff

In Figure 1B, the central sulcus line of subject B is incorrect. The authors have provided a corrected version of Figure 1 here.




Figure 1. Behavioral task and ECoG electrode locations. A) Monkeys performed sequential right arm and hand movements, which consisted of reaching to a knob, grasping the knob with a lateral grip, pulling the knob closer, releasing the knob, and returning the hand to the home position, in a 3-D workspace. During the task, ECoG and EMG signals were recorded simultaneously. B) Schematic diagrams of ECoG electrode locations in left hemisphere. The planar-surface platinum electrode arrays were placed on the gyrus between the central sulcus (CS) and the arcuate sulcus (AS) in the primary motor area. The # indicates the location according to the column of ECoG electrodes. doi:10.1371/journal.pone.0047992.g001

Reference

- Shin D, Watanabe H, Kambara H, Nambu A, Isa T, et al. (2012) Prediction of Muscle Activities from Electrocorticograms in Primary Motor Cortex of Primates. PLoS ONE 7(10): e47992. doi:10.1371/journal.pone.0047992

Citation: The PLOS ONE Staff (2014) Correction: Prediction of Muscle Activities from Electrocorticograms in Primary Motor Cortex of Primates. PLoS ONE 9(3): e92653. doi:10.1371/journal.pone.0092653

Published: March 27, 2014

Copyright: © 2014 The PLOS ONE Staff. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.